ELSEVIER

Volume 564, Numbers 1–2, 14 August 1998

Journal of Organometallic Chemistry

Contents

Palladium(0)-catalyzed regio- and stereoselective addition of heteroatom compounds bearing Si–Se, Ge–Se, and Si–Ge bonds to phenylacetylene
A. Ogawa, H. Kuniyasu, M. Takeba, T. Ikeda, N. Sonoda and T. Hirao 1

Synthesis and structural study of the ate-complex *rac*-$(CH_3)_2C(C_5H_3\text{-}3\text{-}Si(CH_3)_3)_2Yb(\mu_2\text{-}Cl)_2Li(OEt_2)_2$ and the polymeric compound $\{rac\text{-}(CH_3)_2C(C_5H_3\text{-}3\text{-}Si(CH_3)_3)_2Yb[(\mu_2\text{-}H)_2B(\mu_2\text{-}H)_2]_2Li(THF)_2\}_\infty$
A.V. Khvostov, V.K. Belsky, A.I. Sizov, B.M. Bulychev and N.B. Ivchenko 5

Reactions of protic reagents with a tungsten phosphenium ion complex
H.-U. Reisacher, E.N. Duesler and R.T. Paine 13

New osmium hydrides containing the bulky phosphine 1,2-bis(diisopropylphosphino)ethane (dippe)
M. Jiménez Tenorio, M.C. Puerta, I. Salcedo and P. Valerga. 21

Trans-4,5-bis(2-pyridyl)norbornane: a bidentate nitrogen ligand with a potentially large bite angle
J.G. Delis, M. Rep, M. Kranenburg, K. Vrieze, P.W. van Leeuwen, N. Veldman and A.L. Spek. 29

Reversible sulfur dioxide reactions with cyclopentadienylnickel(II) organochalcogenide complexes
J. Darkwa, R.M. Moutloali and T. Nyokong 37

Polymeric organosilicon systems. XXIX. Thermal properties of poly[(disilanylene)oligophenylenes]
J. Ohshita, K. Sugimoto, T. Watanabe, A. Kunai, M. Ishikawa and S. Aoyama 47

Unusual Pt-catalyzed stereo and regioselective intramolecular hydrosilylation of propargyl alcohols to (*E,E*)-1,6-dioxa-2,7-disila-3,8-ecidienes
A.M. Caporusso, S. Barontini, P. Pertici, G. Vitulli and P. Salvadori 57

Stereoselective synthesis of new hetero(P, Si, Ge, Sn)cyclic derivatives from zirconium diyne and diene complexes
M. Mirza-Aghayan, R. Boukherroub, G. Oba, G. Manuel and M. Koenig. 61

Syntheses and properties of tetrathio- and tetraseleno metalates $[(C_5Me_4R)_2NbE_2]_2M$ (E = S, Se; M = Cr, Mo; R = Me, Et) with peripheric niobocene ligands
O. Blacque, H. Brunner, M.M. Kubicki, D. Lucas, W. Meier, Y. Mugnier, B. Nuber, B. Stubenhofer and J. Wachter 71

Zirconocene-catalyzed dehydrogenative coupling of phenylgermane and properties of the resulting partially network polyphenylgermanes
N. Choi and M. Tanaka 81

Synthesis, structure and reactivity of molybdenum imido and oxo complexes. X-ray structure of $MoCl_3(NAr)L_2$ (Ar = 2,6-diisopropyl phenyl; L = N-2',6'-diisopropyl phenyl-2,5-dimethyl benzamide)
F. Javier de la Mata and J.W. Ziller 85

Reactivity of chlorodimethylsilyl-η^5-cyclopentadienyltrichlorotitanium with nitrogen based donors. X-ray molecular structure of $[Ti\{\eta^5\text{-}C_5H_4SiMe_2[\eta^1\text{-}N(2,6\text{-}Me_2C_6H_3)]\}Cl_2]$
R. Gómez, P. Gómez-Sal, A. Martín, A. Núñez, P.A. del Real and P. Royo 93

A synthetic and X-ray crystallographic study of the indenyl-phosphine complexes $1,3\text{-}(Ph_2P{=}X)_2(C_9H_6)$, (X = O, S) and $(\eta^5\text{-}C_9H_5(Ph_2P{=}S)_2)[Mn(CO)_3]$: versatile ligands for the preparation of heteropolymetallic complexes
M. Stradiotto, C.M. Kozak and M.J. McGlinchey 101

Synthesis, characterization and polymerization potential of *ansa*-metallocene dichloride complexes of titanium, zirconium and hafnium containing a Si–N–Si bridging unit
H.G. Alt, K. Föttinger and W. Milius 109

Synthesis, characterization and polymerization potential of unbridged and bridged half-sandwich complexes of zirconium containing *N*-donor ligands
H.G. Alt, K. Föttinger and W. Milius 115

Relationships between basicity, redox behaviour of ferrocenylamines and their reactivity with Pt[II] compounds
N.W. Duffy, J. Harper, P. Ramani, R. Ranatunge-Bandarage, B.H. Robinson and J. Simpson 125

Synthesis and reactivity of tetramethylthioureatriosmium clusters: crystal structures of $[(\mu\text{-H})Os_3(CO)_8(\mu_3\text{-S})(\mu\text{-OMeCO})\{\eta^1\text{-SC}(NMe_2)_2\}]$ and $[Os_3(CO)_9(\mu\text{-OH})(\mu\text{-OMeCO})(PPh_3)]$
S.M. Tareque Abedin, K.A. Azam, M.B. Hursthouse, S.E. Kabir, K.M. Abdul Malik, R. Miah and H. Vahrenkamp 133

Synthesis, structure and reactivity of novel W–Ge chlorocarbonyl compounds. X-ray crystal structure of $[WCl(GeCl_3)(CO)_3(NCEt)_2]$
T. Szymańska-Buzar and T. Głowiak 143

Chirale Bisphosphane VI. Chelatkomplexe von Platin(II) mit racemischen und optisch reinen *trans*-Cyclopentan-1,2-diyl-bis(phosphonigsäurediester)- und *trans*-Cyclopentan-1,2-diyl-bis(phosphonigsäurediamid)-Liganden: Synthesen, Strukturen und C–H-Aktivierungsreaktionen
L. Dahlenburg, C. Becker, J. Höck and S. Mertel 155

Base-free tris(indenyl)lanthanoid(III) complexes (Ln = La, Pr, Nd, Sm): solid-state structure and solution NMR/NIR–vis spectroscopy
J. Guan and R.D. Fischer 167

Halbsandwich-Komplexe von (*S*)-1-Amino-2-(methoxymethyl)-pyrrolidin (SAMP)
W. Hoffmüller, K. Polborn, I. Krossing, H. Nöth and W. Beck 179

[3 + 2] Cycloaddition reactions of the phospha-alkyne, Bu^{t}CP, with a transition metal trisulphide containing terminal thio ligands: syntheses, crystal and molecular structures of $[W(\eta^5\text{-}(C_5Me_5)(S)(S_2PC^tBu)]_2$ and $[W(\eta^5\text{-}(C_5Me_5)(S)(S_2C_2Ph_2)]_2$
S.E. d'Arbeloff, P.B. Hitchcock, J.F. Nixon, T. Nagasawa, H. Kawaguchi and K. Tatsumi 189

Synthesis and structural characterisation of nickel(II) alkyl complexes NiR_2, $[Ni(Cp)R]_2$ and a 'head-to-tail' coupled alkyl intermediate compound [NiR(Cl)(R–R)] [R = $C(SiMe_3)_2C_5H_4N$-2, $CPh(SiMe_3)C_5H_4N$-2 or $CH(SiMe_3)C_5H_4N$-2; Cp = η^5-C_5H_5]
W.-P. Leung, H.-K. Lee, Z.-Y. Zhou and T.C. Mak 193

Reactivity of the 1-azavinylidene cluster $[Ru_3(\mu\text{-H})(\mu\text{-N=CPh}_2)(CO)_{10}]$ with hydrogen, tertiary silanes and tertiary stannanes
C. Bois, J.A. Cabeza, R.J. Franco, V. Riera and E. Saborit 201

New binuclear alkyl and half-sandwich cyclopentadienyl imido titanium complexes containing acetamidinate and benzamidinate supporting ligands
P.J. Stewart, A.J. Blake and P. Mountford 209

Organotin compounds containing a bulky $(Me_3Si)_3C$ or related ligand. Crystal structures of $\{(Me_3Si)_3CMe(O_2NO)Sn\}_2O$, $(PhMe_2Si)_3CSnMeCl_2$ and $(PhMe_2Si)_3CSnCl_3$
S.S. Al-Juaid, M. Al-Rawi, C. Eaborn, P.B. Hitchcock and J.D. Smith 215

Chirale Bisphosphane VIII. Die enantioselektive Hydrierung von α-Acetamidozimtsäure durch kationische Rhodium(I)-Komplexe mit optisch reinen (1*S*,2*S*)-Cyclopentan-1,2-diyl-bis(phosphonigsäurediester)- und (1*S*,2*S*)-Cyclopentan-1,2-diyl-bis(phosphonigsäurediamid)-Liganden
L. Dahlenburg and C. Eckert 227

Transmetallation reaction of cyclomercurated anils of benzoylferrocene with metallic tin. X-ray crystal structure of $[SnCl_2\{(\eta^5\text{-}C_5H_5)Fe(\eta^5\text{-}C_5H_3CPh{=}NC_6H_4\text{-}4\text{-}CH_3)\}_2]$
Y.J. Wu, L. Ding, Z.X. Zhou, C.X. Du and W.L. Wang 233

C–H activation of methyl vinyl ketone in $Ir(acac)\{\eta^2\text{-}CH_2{=}CHC(O)CH_3\}(PCy_3)$
R. Castarlenas, M.A. Esteruelas, M. Martín and L.A. Oro 241

The synthesis of $[AuCo(CO)_4]_2(\mu\text{-dppfe})$ (dppfe = 1,1′-bis(diphenyl phosphino)ferrocene) and the P–Au–Co–CO(axial) skeletal pair formation between neighboring molecules in the crystal
S. Onaka, Y. Katsukawa and M. Yamashita 249

Polynuclear metal complexes linked by conjugated pyridines
J.T. Lin, M.-F. Yang, C. Tsai and Y.S. Wen 257

Synthesis, characterization and chemistry of some long chain alkoxycarbonyl manganese pentacarbonyl complexes $[Mn(CO)_5\{C(O)OR\}]$ (where R = *n*-C_7H_{15} to *n*-$C_{10}H_{21}$, *n*-$C_{12}H_{25}$, *n*-$C_{14}H_{29}$, and *n*-$C_{16}H_{33}$)
X. Yin, J.-A.M. Andersen, A. Cotton and J.R. Moss 267

Synthesis and reactivity of cyclopentadienyl chloro, imido and alkylidene tungsten (VI) complexes
F.J. de la Mata, J. Gómez and P. Royo 277

Book review 283
Author index 285
Subject index 287

The publisher encourages the submission of articles in electronic form thus saving time and avoiding rekeying errors. A leaflet describing our requirements is available from the publisher upon request.